

O VI Column Density Distribution in the Local Bubble - Results from 3D Adaptive Mesh Refinement Simulations

Miguel A. de Avillez

Department of Mathematics, University of Évora, R. Romão Ramalho 59, 7000 Évora, Portugal. Email: mavillez@galaxy.lca.uevora.pt

Dieter Breitschwerdt

Institut für Astronomie, Universität Wien, Türkenschanzstr. 17, A-1180 Wien, Austria. Email: breitschwerdt@astro.univie.ac.at

Abstract. The Local Bubble (LB) is an X-ray emitting region extending 100 pc in radius in the Galactic plane and 300 pc perpendicular to it, and it is embedded in a somewhat larger H I deficient cavity. Its origin and spectral properties in UV, EUV and X-rays are still poorly understood. We have performed 3D high resolution (down to 1.25 pc) hydrodynamic superbubble simulations of the LB and Loop I superbubble in a realistic *inhomogeneous* background ISM, disturbed by supernova (SN) explosions at the Galactic rate. We can reproduce (i) the size of the bubbles (in contrast to similarity solutions), (ii) the interaction shell with Loop I, discovered with ROSAT, (iii) predict the merging of the two bubbles in about 3 Myr, when the interaction shell starts to fragment, and, (iv) the generation of blobs like the Local Cloud as a consequence of a dynamical instability. The O VI column densities are monitored and found to be in excellent agreement with COPERNICUS and FUSE absorption line data, showing LB column densities $< 1.7 \times 10^{13} \text{ cm}^{-2}$, in contrast to other existing models.

1. Introduction

Standard Local Bubble (LB) models fail to reproduce the observed low O VI absorption column density (Shelton & Cox 1994; for a recent discussion see Breitschwerdt & Cox 2004). Heliospheric in situ measurements are sensitive to the boundary conditions imposed by the Local Bubble and the O VI column density in absorption is a crucial test for modelling of the local ISM (Cox 2004). It seems most plausible that the LB is the result of 20 successive explosions, originating from stars with masses between 11 and 20 $M_{\odot}$ from the moving subgroup B1 of the Pleiades in the last 14.5 Myrs (Berghöfer & Breitschwerdt 2002). We have simulated the LB evolution by means of a 3D AMR hydrocode and calculated the observed O VI column density in absorption (along lines of sight crossing Loop I).

2. Model and Simulations

We use the 3D model of Avillez (2000), where the ISM is disturbed by supernova (SN) explosions at the Galactic rate, and took data cubes of previous runs with

a finest adaptive mesh refinement resolution of 1.25 pc (Avillez & Breitschwerdt 2004). We then picked up a site with enough mass to form the 81 stars, with masses, M_* , between 7 and 31 $M_\odot$, that compose the Sco Cen cluster; 39 massive stars with $14 \leq M_* \leq 31 M_\odot$ have already gone off, generating the Loop I cavity.

Jpeg image LB.jpg

Presently the Sco Cen cluster (here located at (375, 400) pc) has 42 stars to explode in the next 13 Myrs). We followed the trajectory of the moving subgroup B1 of Pleiades, whose SNe in the LB went off along a path crossing the solar neighbourhood (Fig. 1). Periodic boundary conditions are applied along the four vertical boundary faces, while outflow boundary conditions are imposed at the top ($z = 10$ kpc) and bottom ($z = -10$ kpc) boundaries. The simulation time of this run was 30 Myr.

Figure 1. O VI contour map of a 3D Local Bubble simulation 14 Myr after the first explosion; LB is centered at (175, 400) pc and Loop I at (375, 400) pc.

3. Results

Morphology: The locally enhanced SN rates produce coherent LB and Loop I structures (due to ongoing star formation) within a highly disturbed background medium. The successive explosions heat and pressurize the LB, which at first looks smooth, but develops internal temperature and density structure at later stages. After 14 Myr the LB cavity, bounded by an outer shell, which will start to fragment due to Rayleigh-Taylor instabilities in ~ 3 Myr from now, fills a volume roughly corresponding to the present day size (Fig. 1).




Figure 2. O VI column density averaged over angles (left panel) indicated in Fig. 2. and maximum column density (right panel) as a function of LOS path length at $14.1 \leq t \leq 15$ Myr of Local and Loop I bubbles evolution.

Oxygen VI: The average and maximum column densities of O VI, i.e., $\langle N(\text{O VI}) \rangle$ and $N_{\text{max}}(\text{O VI})$ are calculated along 91 lines of sight (LOS) ex-

tending from the Sun and crossing Loop I from an angle of -45° to $+45^\circ$ (s. Fig. 1). Within the LB (i.e., for a LOS length $l_{LOS} \leq 100$ pc) $\langle N(\text{O VI}) \rangle$ and $N_{\text{max}}(\text{O VI})$ decrease steeply from 5×10^{13} to $3 \times 10^{11} \text{ cm}^{-2}$ and from 1.2×10^{14} to $1.5 \times 10^{12} \text{ cm}^{-2}$, respectively, for $14.1 \leq t \leq 15$ Myr (Fig. 2), because no further SN explosions occur and recombination is taking place. For LOS sampling gas from outside the LB (i.e., $l_{LOS} > 100$ pc) $\langle N(\text{O VI}) \rangle > 6 \times 10^{12}$ and $N_{\text{max}}(\text{O VI}) > 5 \times 10^{13} \text{ cm}^{-2}$. The histograms of column densities obtained in the 91 LOS for $t = 14.5$ and 14.6 Myr (Fig. 3) show that for $t = 14.6$ Myr all the LOS have column densities smaller than $10^{12.9} \text{ cm}^{-2}$, while for $t = 14.5$ Myr 67% of the lines have column densities smaller than 10^{13} cm^{-2} and in particular 49% of the lines have $N(\text{O VI}) \leq 7.9 \times 10^{12} \text{ cm}^{-2}$.




Figure 3. Histogram of the percentage of LOS with various ranges of observed $N(\text{O VI})$ within the LB at $t = 14.5$ and 14.6 Myr.

Myr ago, whereas the nearest explosions to the Sun happened 2.8 ± 0.7 Myr ago, in good agreement with the most recent dating inferred from ^{60}Fe isotope analysis in the ferromanganese crust of deep ocean layers (Knie et al. 2004).

Acknowledgements M.A. would like to thank the organization for the financial support to attend this excellent conference.

References

- Avillez, M. A. 2000, MNRAS, 315, 479.
- Avillez, M. A., & Breitschwerdt, D. 2004, A&A, 425, 899
- Berghöfer, T., & Breitschwerdt, D. 2002, A&A, 390, 299.
- Breitschwerdt, D., & Cox, D. P. 2004, in “How does the Galaxy Work?”, eds. E. Alfaro, E. Perez, & J. Franco, Kluwer (Dordrecht), p. 391
- Cox, D. P. 2004, Ap&SS, 289, 469
- Knie, K., et al. 2004, Phys. Rev. Lett., 93, 17
- Oegerle, W. R., et al. 2004, ApJ, submitted [astro-ph/0411065]
- Shelton, R., & Cox, D. P. 1994, ApJ, 434, 599

Noting that in the present model for $t \geq 14.5$ Myr the O VI column densities are smaller than $1.7 \times 10^{13} \text{ cm}^{-2}$ and $\langle N(\text{O VI}) \rangle \leq 8.5 \times 10^{12} \text{ cm}^{-2}$ (blue line in both panels of Fig. 2) and that a mean column density of $7 \times 10^{12} \text{ cm}^{-2}$ is inferred from analysis of FUSE absorption line data in the Local ISM (Oegerle et al. 2004) we estimate that the present LB time is $14.7_{-0.2}^{+0.5}$ Myr.

4. Conclusions

The O VI column density is a sensitive tracer of the age of an *evolved* superbubble, and can thus give a constraint on the timescale of the last explosion, which occurred in the present simulations $0.6_{-0.2}^{+0.5}$

This figure "LB.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0501466v2>